

Could we learn more about HERA high Q^2 anomaly from LEP200 and TEVATRON ?

A.S.Belyaev ^{a,1}, A.V.Gladyshev ^{b,2}

^a *Skobeltsin Institute for Nuclear Physics, Moscow State University,
119 899, Moscow, RUSSIA*

^b *Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,
141 980 Dubna, Moscow Region, RUSSIA*

Abstract

The excess of high Q^2 events at HERA has been the subject of recent extensive studies in the framework of several models related to new physics. Here, we would like to concentrate on the most promising, from our point of view, model describing HERA anomaly. We investigate HERA events within the R -parity broken SUSY model and check its relation to LEP and TEVATRON colliders. This study shows that if a squark resonance really takes place at HERA, supersymmetry with broken R -parity can be revealed at either LEP200 or TEVATRON.

¹e-mail: belyaev@monet.npi.msu.su

²e-mail: gladyshev@thsun1.jinr.dubna.su

1 Introduction

Anomalous high- Q^2 events recently observed at HERA by the H1 [1] and ZEUS [2] collaborations have provoked much efforts of their explanation within different theoretical frameworks [3, 4]. The proposed explanations are contact interactions, leptoquarks and signals of R -parity violation in supersymmetric interactions. The contact interactions do not give us satisfactory explanation of either the shape of Q^2 distribution of HERA events or the distribution over the $e + jet$ invariant mass which has a peak around 200 GeV. We would like to take a fresh look at the possibility of R -parity violation providing a s -channel resonance at HERA.

The starting point for our analysis is the superpotential of the Minimal Supersymmetric Standard Model with R -violating terms included

$$W = h_E L E^c H_1 + h_D Q D^c H_1 + h_U Q U^c H_2 + \mu H_1 H_2 \\ + \mu_i H_2 L_i + \lambda_{ijk} L_i L_j E_k^c + \lambda'_{ijk} L_i Q_j D_k^c + \lambda''_{ijk} U_i^c D_j^c D_k^c \quad (1)$$

The terms with λ and λ' violate a lepton number while the last term with λ'' violates a baryon number. B and L operators, if they exist simultaneously, lead to a fast proton decay [8]. Thus, only one kind of operators is allowed. HERA events can be explained in the framework of the R -parity broken SUSY model only if $\lambda' \neq 0$. That is why we hereafter consider the case when $\lambda'' = 0$.

The λ and λ' -terms result in the following lagrangian for component fields:

$$L_{R_p, \lambda} = \lambda_{ijk} [\tilde{\nu}_{iL} \bar{e}_{kR} e_{jL} + \tilde{e}_{jL} \bar{e}_{kR} \nu_{iL} + \tilde{e}_{kL}^* (\overline{\nu_{iL}})^C e_{jL} - (i \leftrightarrow j)]_F + h.c., \quad (2)$$

$$L_{R_p, \lambda'} = \lambda'_{ijk} [\tilde{\nu}_{iL} \bar{d}_{kR} d_{jL} + \tilde{d}_{jL} \bar{d}_{kR} \nu_{iL} + \tilde{d}_{kL}^* (\overline{\nu_{iL}})^C d_{jL} - \\ \tilde{e}_{iL} \bar{d}_{kR} u_{jL} - \tilde{u}_{jL} \bar{d}_{kR} e_{iL} - \tilde{d}_{kL}^* (\overline{e_{iL}})^C u_{jL}] + h.c. \quad (3)$$

The present experimental limits on different couplings are presented in table 1.

$\lambda_{13k} \leq 0.2$	(a)[6]	$\lambda'_{111} \leq 7 \cdot 10^{-3}$	(c)[7]	$\lambda'_{131} \leq 3.7 \cdot 10^{-2}$	(b)[8]	$\lambda'_{23k} \leq 0.24$	(a)[6]
$\lambda_{23k} \leq 0.2$	(a)[6]	$\lambda'_{ijk} \leq 0.024$	(d)[9]	$\lambda'_{132} \leq 7.8 \cdot 10^{-3}$	(b)[8]	$\lambda'_{ijk} \leq 0.52$	(e)[10]
$\lambda_{12k} \leq 0.08$	(a)[6]			$\lambda'_{133} \leq 1.7 \cdot 10^{-3}$	(b)[8]		

Table 1: The strictest experimental bounds on R -parity violating couplings. Limits derived from the following processes: a) charged-current universality, b) ν_e mass; c) neutrinoless double beta-decay; d) $BR(K^+ \rightarrow \pi^+ \nu \bar{\nu})$; e) atomic parity violation and eD asymmetry.

Bounds shown in table 1 scale with the following factors:

a), d) and e) – $M_{\tilde{q}}/200 \text{ GeV}$;

b) – $\sqrt{M_{\tilde{q}}/200 \text{ GeV}}$;

c) – $(M_{\tilde{q}}/200 \text{ GeV})^2 \sqrt{M_{\tilde{g}}/1 \text{ TeV}}$

The purpose of the present letter is to re-examine the possible explanation of the excess of high Q^2 neutral current DIS events at HERA assuming R -parity violation and discuss its possible check at LEP200 and TEVATRON.

It should be stressed that all analytical and the most part of numerical calculations have been made by the CompHEP package [11]. This package allows one to perform the complete tree level calculations in the framework of any fed model. We have implemented into this program the part of supersymmetric standard model with R-parity violating terms relevant for our analysis.




Figure 1: Diagrams for $e^+ + p \rightarrow e^+ jet + X$ HERA events in the R -parity violating SUSY model.




Figure 2: Q^2 distribution, predicted by SM (solid line), SM+ R -parity breaking squark resonance (dashed line) [$\lambda' = 0.047$] and data (solid dots).

2 R -parity violation at HERA

To describe HERA anomalous events, we used data obtained by these collaborations when 10 events survived after the $Q^2 > 20000 \text{ GeV}^2$ cut, while only 4 events were expected according to Standard Model predictions. We also took into account the integrated luminosity ZEUS+H1 equal to 34.4 pb^{-1} and an average detector efficiency 0.8.

In our calculations we used the CTEQ3M structure function [13] and Q^2 scale $(p_e - p'_e)^2$. The total cross section describing the observed data turned out to be 0.36 pb, while the Standard Model gives only 0.16 pb. Five events observed by the H1 collaboration ($Q^2 > 20000 \text{ GeV}^2$) have a clear peak for $m_{ej} = 200 \text{ GeV}$. ZEUS events are more spread over m_{ej} , but also have a mean value around 200 GeV. That is why the natural mass of sparticles describing these events is 200 GeV.

We have calculated the complete set of Feynman diagrams of the process $e^+ + p \rightarrow e^+ + jet + X$ in the framework of the R -broken supersymmetric model for HERA energy. The relevant graphs are shown in fig.1. As is naturally expected the main contribution comes from diagrams with a resonant squark in the s -channel (99%). Contributions from the t -channel diagrams are negligible (1%) for the case under consideration. However, these diagrams give the same order contribution as a *nonresonant* s -channel diagram with a heavy sparticle, but this is not the subject of our study.

We use values for R -parity violating Yukawa couplings, which satisfy existing limits, see table 1. We assumed all couplings non-zero. Couplings λ'_{111} , λ'_{131} , λ'_{132} and λ'_{133} were fixed at their upper bounds coming from the electron neutrino mass limits. The rest λ 's were put equal to each other and varied in the allowed range. The presence of non-zero couplings implies the branching ratio $BR(\tilde{q} \rightarrow e^+ q)$ to be smaller than unity, even if the R -parity conserving decay channel is suppressed. For the values of λ 's under consideration we have $BR(\tilde{t} \rightarrow e^+ q) \simeq 2/7$ and $BR(\tilde{c} \rightarrow e^+ q) \simeq 1/3$. Due to the smallness of λ'_{111} the diagram with the $\tilde{u}$ exchange is neglected hereafter.

We obtained the value of λ' equal to 0.047 which describes the event excess in HERA data with the 200 GeV s -channel squark resonance. This value is higher than the 0.024 limit coming from [9]. But the limit on coupling in [9] was obtained under the assumption that only one λ was non-zero, which is not the case of this study, and as it was shown in [4] this limit can be relaxed.

Figure 2 shows the Q^2 distribution for the Standard Model and SUSY model with broken R -parity tuned to describe data, and combined ZEUS+H1 data distribution itself.

One can see that the shape of deviation from SM is perfectly described by the s -channel squark resonance.

3 Implication for LEP200 and TEVATRON

Having in mind that processes with R -parity violation could occur in e^+e^- collisions at LEP, we consider the possibility of detecting it in the pair $b\bar{b}$ production via sparticle exchange. Comparing to the Standard Model, some excess in dijet production is expected. For this purpose, we have calculated the complete set of relevant diagrams presented in fig.3. Hereafter, we assume that sleptons have mass 200 GeV if not mentioned specially.

There is a big difference in the up and down quark production. There is an additional s -channel diagram with sneutrino for the down quark production, which makes the total

cross section dependent not only on λ' but also on λ values. If we put λ equal to its upper limit (0.2), see table 1, then we will have 25% deviation $\simeq 1.1$ pb of the total cross section from the Standard Model for $\sqrt{s} = 180$ GeV LEP. In our calculation we put the value of λ' to those describing HERA data — 0.047. The dependence of the total $q_d\bar{q}_d$ cross section on λ' is shown in fig.4a). The value of λ of an order of 0.1-0.2 leads to the 15-25% excess in the $b\bar{b}$ production cross section. This deviation from the SM R_b prediction can be measured at LEP. As for the up quark production, t -channel diagrams bring a very small negative contribution for values of λ' s under consideration. The dependence of the total $q_u\bar{q}_u$ cross section as a function of λ' is shown in fig.4b). One could expect some contribution of a 200 GeV SUSY particle to the very accurately measured R_b value for LEP90 which is equal to 0.2178 ± 0.0011 and still has 1.5σ deviation from theoretical predictions. But it turned out that for $\sqrt{s} = 90$ GeV the contribution from additional diagrams to the $b\bar{b}$ production was only 0.021 pb, while the total production $b\bar{b}$ rate was 9040 pb. Thus, this additional SUSY contribution changes R_b only in the 6-th digit.




Figure 3: Diagrams for the $e^+e^- \rightarrow q\bar{q}$ process in the R -parity violated SUSY model




Figure 4: Total cross section of $e^+e^- \rightarrow q\bar{q}$ versus λ' for down (a) and up (b) quark pair production at LEP. Dashed line — SM prediction.

Though limits on λ 's are not so strict, it seems unnatural if their values are much higher than λ' . So we would like to look at the case when λ is small or equal to 0 and the excess of the $b\bar{b}$ production is absent. For example, for $\lambda \simeq \lambda' = 0.047$ the $b\bar{b}$ production deviates from the Standard Model only by 5%. By a lucky chance we have at present three different colliders with energy of an order of 1 TeV: lepton-lepton — LEP, lepton-hadron — HERA and hadron-hadron — TEVATRON. These machines are highly complementary to each other. We show below that possible failure to find R -parity broken SUSY (if it really exists) at LEP leads to the luck at TEVATRON in this search and vice versa.

For TEVATRON one of the the most interesting effects of revealing R -parity violation can be the resonance single top production via a slepton decay.

The complete set of Feynman diagrams for the $2 \rightarrow 2$ process of single top production is shown in fig.5.




Figure 5: Diagrams for $2 \rightarrow 2$ single top production processes at TEVATRON in the R -parity violated SUSY model.




Figure 6: Total cross section of the $p\bar{p} \rightarrow t\bar{q}_d(\bar{t}q_d)$ single top production process versus λ (a) and total cross section of the $b\bar{b}$ production at LEP180 versus λ (b). Dashed line shows the SM prediction.

In spite of that the cross section is directly proportional only to λ' , there is also dependence of the cross section on the λ coupling which is directly related to the width

and branching ratios of the selectron decay. For comparatively big λ (0.1-0.2) the selectron mainly decays leptonically and the single top production is suppressed. In this case, the deviation in the $b\bar{b}$ production could be measured at LEP. On the other hand, if λ is small or even equal to 0, there is a significant contribution to the single top production via the resonant selectron diagram. This contribution can be as large as 100% of the total single top production rate of SM $q\bar{q}' \rightarrow t\bar{q}_d$. It is clearly illustrated in fig 6 where the dependence of this deviation versus λ (λ' is fixed to 0.047) is shown. If we assume the total cross section of the single top $t\bar{b}$ ($+\bar{t}b$) production including NLO corrections [12] (we use NLO K-factor 1.5) to be about 0.9 pb, for $\lambda=0.0-0.1$ the deviation from the SM prediction is 50–100%. The same effect is expected for the $W - gluon$ fusion single top production process which is a separate subject of study and not discussed here. We can see that by lucky coincidence LEP and TEVATRON magically complement each other in search for supersymmetry with broken R -parity, which naturally describes high Q^2 HERA events.

Conclusions

We have demonstrated that HERA events can be explained within the Supersymmetric Standard Model with broken R -parity. The Yukawa couplings responsible for processes under study do not exceed their experimental bounds. Processes with R -parity breaking can also take place in e^+e^- collisions at LEP200 and TEVATRON, which are complementary to each other. If R -parity violation really takes place, as has been shown above, it can be revealed at either TEVATRON or LEP in the near future.

We would like to thank D.I.Kazakov and W. de Boer for valuable discussions and stimulation of this study. Our work has been supported by Russian Foundation for Basic Research, grants # 96-02-17379, #96-02-19773-a and ICFPM 1996 grant.

References

- [1] G.Adolf et al., H1 collaboration, DESY preprint 97-24, [hep-ex/9702012](#).
- [2] J.Breitweg et al., ZEUS collaboration, DESY preprint 97-25, [hep-ex/9702015](#).
- [3] S. L. Adler, [hep-ph/9702378](#);
D. Choudhury, S.Raychaudhury, [hep-ph/9702392](#);
T. K. Kuo, T. Lee, [hep-ph/9703255](#);
H. Dreiner, P. Morawitz, [hep-ph/9703279](#);
J. Blumlein, [hep-ph/9703287](#);
J. Kalinowski, R. Ruckl, H. Spiesberger, P.M.Zerwas, [hep-ph/9703288](#);
K. S. Babu, C. Kolda, J. March-Russell, F. Wilczek, [hep-ph/9703299](#);
- [4] G. Altarelli, J. Ellis, G. F. Guidice, S. Lola, M. L. Mangano, [hep-ph/9703276](#);
- [5] A. Bouquet and P. Salati, Nucl. Phys. **B284** (1987) 557;
B.A. Campabell, S. Davson, J. Ellis and K. Olive, Phys. Lett. **B256** (1991) 457;

- A. Nelson and S.Barr, Phys. Lett. **B246** (1990) 141
- [6] V.Barger, G.F. Giudice and T.Han, Phys. Rev. D 40 (1989) 2887
 - [7] M. Hirsh, H.V. Klapdor-Kleingrothaus and S.G. Kovalenko, Phys. Rev. Lett. **75** (1995) 17; Phys. Rev. **D53** (1996) 1329
 - [8] R.M. Godbole, P. Roy, and X. Tata, Nucl. Phys. **B401**(1993) 67
 - [9] H. Agashe and M. Graesser, Phys. Rev. **D54**(1995) 4445
 - [10] G. Bhattacharyya, J. Ellis and K. Sridhar, Mod. Phys. Lett. **A10** (1995) 1583
 - [11] E.E.Boos et al., hep-ph/9503280,SNUTP-94-116;
E.E.Boos et al., in Proc. of the Xth Int. Workshop on High Energy Physics and Quantum Field Theory, QFTHEP-95, ed. by B.Levtchenko, and V.Savrin, (Moscow, 1995), p.101
 - [12] Martin C. Smith, S. Willenbrock, Phys.Rev. **D54** (1996) 6696
 - [13] CTEQ Collaboration: H.L. Lai et al., Phys. Rev. **D51** , (1995) 4763

